

# Population health profile of the Hunter Urban Division of General Practice

Population Profile Series: No. 16

PHIDU

November 2005



PHIDU



Australian Government

Australian Institute of  
Health and Welfare



THE UNIVERSITY  
OF ADELAIDE  
AUSTRALIA

SUB CRUCE LUMEN

© Commonwealth of Australia 2005

This work may be reproduced and used subject to acknowledgement of the source of any material so reproduced.

## National Library of Australia Cataloguing in Publication entry

---

Population health profile of the Hunter Urban Division of General Practice.

Bibliography.

ISBN 0 7308 9423 1.

1. Public health - New South Wales - Hunter - Statistics. 2. Health status indicators - New South Wales - Hunter - Statistics. 3. Health service areas - New South Wales - Hunter. 4. Hunter (N.S.W.) - Statistics, Medical. I. Public Health Information Development Unit (Australia). II. Australia. Dept. of Health and Ageing. III. Australian Institute of Health and Welfare. (Series : Population profile series, 1833-0452 ; no. 16).

362.1099441

ISSN 1833-0452 Population Profile Series

**Public Health Information Development Unit, The University of Adelaide**  
***A Collaborating Unit of the Australian Institute of Health and Welfare***

---

This profile was produced by PHIDU, the Public Health Information Development Unit at The University of Adelaide, South Australia. The work was funded under a grant from the Australian Government Department of Health and Ageing. The views expressed in this profile are solely those of the authors and should not be attributed to the Department of Health and Ageing or the Minister for Health and Ageing.

The data in this report are designed to be used for needs assessment and planning purposes: while they are based on the best available data and analytic processes, data available by postcode or Statistical Local Area, as used in this report, cannot be precisely translated to Division. Division totals in the report should, therefore, be seen as estimates. Interpretation of differences between data in this profile and similar data from other sources needs to be undertaken with care as such differences may be due to the use of different methodology to produce the data.

Suggested citation:

PHIDU. (2005) *Population health profile of the Hunter Urban Division of General Practice*. Population Profile Series: No. 16. Public Health Information Development Unit (PHIDU), Adelaide.

Enquiries about or comments on this publication should be addressed to:

PHIDU, The University of Adelaide, South Australia 5005  
Phone: 08-8303 6237 or e-mail: [PHIDU@publichealth.gov.au](mailto:PHIDU@publichealth.gov.au)

This publication, the maps and supporting data, together with other publications on population health, are available from the PHIDU website ([www.publichealth.gov.au](http://www.publichealth.gov.au)).

Published by Public Health Information Development Unit, The University of Adelaide

Contributors: Anthea Page, Sarah Ambrose, Liz Fisher, Kristin Leahy and John Glover

# Population health profile

## of the *Hunter Urban Division of General Practice*

### Introduction

This profile has been designed to provide a description of the population of the Hunter Urban Division of General Practice, and aspects of their health. Its purpose is to provide information to support a population health approach, which aims to improve the health of the entire population and to reduce health inequalities among population groups: a more detailed discussion of a population health approach is provided in the supporting information, page 17.

### Contents

The profile includes a number of tables, maps and graphs to profile population health in the Division and provides comparisons with other areas (eg. country New South Wales and Australia). Specific topics covered include:

- a socio-demographic profile (pages 2-6);
- GP workforce data (page 7);
- immunisation rates (page 7);
- rates of premature death (page 8); and
- estimates of the prevalence of chronic disease and selected risk factors (pages 9-13).

### Key indicators

**Location:** New South Wales

**Division number:** 217

<b>Population‡:</b>	<b>No.</b>	<b>%</b>
Total	443,089	
65+	68,306	15.4%
<25	146,195	33.0%
Indigenous	9,007	2.1%

**Disadvantage score<sup>1</sup>:** 975

**GP services per head of population:**

Division‡	4.3
Australia	4.7

**Population per FTE GP:**

Division‡	1,551
Australia	1,403

**Premature death rate<sup>2</sup>:**

Division‡	328.4
Australia	290.4


<sup>1</sup> Numbers below 1000 (the index score for Australia) indicate the Division is relatively disadvantaged

<sup>2</sup> Deaths at ages 0 to 74 years per 100,000 population


<sup>3</sup> ‡ See note "Data converters and mapping" re calculation of Division Total

## Hunter Urban Division of General Practice

*NSW Divisions of General Practice*



*Hunter Urban DGP by SLA*




# Socio-demographic profile

## Population


The Hunter Urban DGP had an Estimated Resident Population of 443,089 at 30 June 2004.

**Figure 1: Annual population change, Hunter Urban DGP‡, country New South Wales<sup>1</sup>, New South Wales and Australia, 1991 to 1996, 1996 to 2001 and 2001 to 2004**



Over the five years from 1991 to 1996, the Division's population increased by 0.8% on average each year, lower than for country New South Wales (0.9%) and New South Wales (1.0%) and Australia as a whole (1.2%). From 1996 to 2001, the annual percentage increase in the Division of 1.3% was higher than for country New South Wales (1.1%) and equal to New South Wales. The lower growth rate (0.7% per year) from 2001 to 2004 was higher than for country New South Wales and New South Wales (0.6%).

**Figure 2: Population in Hunter Urban DGP‡ and Australia, by age and sex, 2003**



The age distribution of the Division's population is similar to that for Australia overall. The most notable differences are:

- at the youngest ages – a marginally lower proportion of children aged 0 to 4 years;
- from 25 to 44 years – lower proportions of both males and females; and
- at older ages – slightly higher proportions of both males aged 60 years and over, and females aged 55 years and over.

**Table 1: Population by age, Hunter Urban DGP‡ and Australia, 2004**

Age group (years)	Hunter Urban DGP		Australia	
	No.	%	No.	%
0-14	86,108	19.4	3,978,751	19.8
15-24	60,087	13.6	2,762,769	13.8
25-44	119,697	27.0	5,881,048	29.3
45-64	108,891	24.6	4,864,037	24.2
65-74	34,691	7.8	1,374,792	6.8
75-84	25,747	5.8	934,505	4.7
85+	7,868	1.8	295,602	1.5
<b>Total</b>	<b>443,089</b>	<b>100.0</b>	<b>20,091,504</b>	<b>100.0</b>

As shown in the age-sex pyramid above, Hunter Urban DGP had a lower proportion of the population aged 25 to 44 years (27.0%) compared to Australia (29.3%) (Table 1). Conversely, the 65 years and over age groups had higher proportions compared to Australia as a whole.

The Hunter Urban DGP comprised 4.0% of people born in predominantly non-English speaking countries and resident in Australia for five years or more (Table 2), similar to the 4.1% in country New South Wales. Recent arrivals (those resident in Australia for less than five years) from non-English speaking countries comprised 0.6% of the Division's population (consistent with the 0.5% in country New South Wales).

<sup>1</sup>References to 'country New South Wales' relate to New South Wales excluding the Sydney Statistical Division.

‡ See note under 'Data converters and mapping' re calculation of Division totals on this page


Of these residents, 0.6% had poor proficiency in English (determined when people aged five years and over born overseas in predominantly non-English speaking countries reported in the Census speaking another language and speaking English 'not well' or 'not at all'), the same as the proportion in country New South Wales, but notably less than New South Wales (3.2%) and Australia (2.4%).

**Table 2: Non-English speaking born, Hunter Urban DGP, country NSW, New South Wales and Australia, 2001**

People born in predominantly non-English speaking countries	Hunter Urban DGP		Country NSW		New South Wales		Australia	
	No.	%	No.	%	No.	%	No.	%
Resident in Australia for five years or more	16,700	4.0	97,983	4.1	803,824	12.7	2,019,410	10.8
Resident in Australia for less than five years	2,359	0.6	12,392	0.5	182,972	2.9	408,074	2.2
Poor proficiency in English <sup>1</sup>	2,218	0.6	13,587	0.6	189,874	3.2	425,399	2.4

<sup>1</sup> Calculated on persons aged 5 years and over who reported speaking another language and speaking English 'not well' or 'not at all'

**Figure 3: Major non-English speaking birthplaces, Hunter Urban DGP, 2001**



## Socioeconomic status

*The indicators presented in this section describe geographic variations in the distribution of the population for a number of key socioeconomic influences, which impact on the health and wellbeing of populations.*

The Hunter Urban DGP had a similar proportion of single parent families (11.9%) to country New South Wales as a whole (11.7%), but fewer Aboriginal and Torres Strait Islanders (2.1%, compared with 3.7% for country New South Wales) (Figure 4, Table 3).

Full-time secondary school education participation of 16 year olds living in the Division (73.3%) was similar to that for country New South Wales (73.4%).

A similar proportion of the Division's households received rent assistance from Centrelink (18.0%) compared to country New South Wales (18.3%), but there were more dwellings rented from the State housing authority (5.8%, compared to 4.6%). The proportion of dwellings with no access to a motor vehicle (11.8%) was higher than that for country New South Wales (10.2%).


The Division had higher proportions of the population who reported using, at home, a computer (40.3%) and the Internet (25.0%), compared to country New South Wales (37.0% and 22.2%).

These socioeconomic indicators show the Division to comprise a population of average socioeconomic status: see also the note on page 5 (Summary of socioeconomic ranking).


**Figure 4: Socio-demographic indicators, Hunter Urban DGP, country NSW, New South Wales and Australia, 2001**

*Note the different scales*


**Single parent families**




**Indigenous‡**




**Education participation at age 16‡**




**Unemployment rate (June 2003)‡**




**Households receiving rent assistance & Dwellings rented from State housing authority**




**Dwellings with no motor vehicle**



**Computer use at home**



**Internet use at home**



‡ See note under 'Data converters and mapping' re calculation of Division totals

**Table 3: Socio-demographic indicators, Hunter Urban DGP, country New South Wales, New South Wales and Australia, 2001**

Indicator	Hunter Urban		Country NSW		NSW		Australia	
	No.	%	No.	%	No.	%	No.	%
Single parent families	13,389	11.9	73,805	11.7	172,199	10.4	529,969	10.7
Indigenous‡	9,007	2.1	91,036	3.7	134,886	2.1	458,261	2.4
Full-time secondary school education at age 16‡	4,289	73.3	24,254	73.4	65,205	75.2	130,198	78.7
Households: rent assistance	28,041	18.0	156,074	18.3	343,540	15.5	1,006,599	15.0
Dwellings rented from the State housing authority	9,265	5.8	41,406	4.6	114,130	4.9	317,171	4.5
Dwellings: no motor vehicle	18,817	11.8	92,576	10.2	280,434	12.0	708,073	10.0
Computer use at home	165,644	40.3	874,207	37.0	2,600,257	41.2	7,881,983	42.0
Internet use at home	103,965	25.0	523,994	22.2	1,751,626	27.7	2,019,410	27.7

‡ See note under 'Data converters and mapping' re calculation of Division total

The unemployment rate of 7.3% in Hunter Urban DGP was the same as country New South Wales and higher than New South Wales (6.0%) (Figure 4, Table 4). The labour force participation rate (71.2%) was lower than country New South Wales (72.3%) and New South Wales (74.6%), while the female labour force participation rate (67.3%) was marginally higher than country New South Wales (66.8%), but lower than New South Wales (69.0%).

**Table 4: Unemployment and labour force, Hunter Urban DGP, country New South Wales, New South Wales and Australia, 2003**

Labour force indicators	Hunter Urban		Country NSW		NSW		Australia	
	No.	%	No.	%	No.	%	No.	%
Unemployment rate ‡	14,910	7.3	83,231	7.3	198,946	6.0	623,791	6.2
Labour force participation‡	204,258	71.2	1,142,496	72.3	3,331,064	74.6	10,038,147	75.2
Female labour force participation (2001)	67,431	67.3	361,345	66.8	1,093,243	69.0	3,306,521	69.7

‡ See note under 'Data converters and mapping' re calculation of Division total


## Summary of the socioeconomic ranking of the Hunter Urban DGP

Following the 2001 Census, the Australian Bureau of Statistics (ABS) produced four socioeconomic indexes for areas (SEIFA) which describe various aspects of the socioeconomic profile of populations in areas. The scores for these indexes for each Statistical Local Area (SLA) or part SLA in Hunter Urban DGP are shown in the supporting information in Table 9, page 17: SLAs are described on page 18.

The Hunter Urban DGP area's SEIFA Index of Relative Socio-Economic Disadvantage (IRSD) score is 975, marginally (2.5%) below the average score for Australia (1000), but above country New South Wales (973); this highlights the near-average socioeconomic status profile of the Hunter Urban DGP population. Variations in the IRSD within the Division are shown in Map 1 at the SLA level.

**Map 1: Index of Relative Socio-Economic Disadvantage by SLA, Hunter Urban DGP, 2001**

*See note under 'Methods' re Data converters and mapping concerning SLAs mapped to the Division. This is of particular relevance where part of an SLA is mapped to the Division.*



## General medical practitioner (GP) supply

A total of 284.9 full-time equivalent (FTE) GPs and 339.1 full-time workload equivalent (FWE<sup>2</sup>) GPs worked in the Division in 2003/04 (Table 5). Of the FWE GPs, 28.0% were female, and 36.1% were over 55 years of age (compared to 26.4% and 33.4%, respectively, for New South Wales).

Apart from the estimated day-time population, the rates of population per FTE GP varied, depending on the population measure used, from a high of 1,551 people per GP (calculated on the average Estimated Resident Population (ERP) as at 30 June 2003 and 30 June 2004), to a low of 1,470 per GP (calculated on the 1 August 2001 Census count – all people counted in the Division on Census night, including visitors from Australia and overseas). The rates of population per FWE were lower, ranging from 1,235 (calculated on the Census count) to 1,303 (calculated on the ERP). When calculated on the estimated daytime population, the rates of population per GP in the Division were 4.3% below those calculated on the Usual Resident Population (usual residents of the Division counted in Australia on Census night).

Based on the ERP, the rates of population per GP in Hunter Urban DGP were higher than the rates for New South Wales and Australia, indicating a lower level of provision of GP services in the Division.

**Table 5: Population per GP in Hunter Urban DGP, New South Wales and Australia, 2003/04**

Population measure	Population	GPs		Population per GP	
		FTE	FWE	FTE	FWE
<b>Hunter Urban DGP</b>					
Census count (adjusted)*	418,786	284.9	339.1	1,470	1,235
Usual Resident Population (URP) (adjusted)*	423,084	..	..	1,485	1,248
Estimated Resident Population (ERP)	441,862	..	..	1,551	1,303
Day-time population (estimated on URP)* ‡	404,859	..	..	1,421	1,194
<b>New South Wales (ERP)</b>	<b>6,706,674</b>	<b>4,819</b>	<b>5,969</b>	<b>1,392</b>	<b>1,124</b>
<b>Australia (ERP)</b>	<b>19,989,303</b>	<b>14,246</b>	<b>16,872</b>	<b>1,403</b>	<b>1,185</b>

\* The Census count, Usual Resident Population and Daytime population were adjusted to reflect population change between 2001 and 2003/2004, as measured by the ERP

‡ See note under 'Data converters and mapping' re calculation of Division totals

## Immunisation

Data from the Australian Childhood Immunisation Register show that 95.9% of children in the Division in 2002 were fully immunised at age one, marginally above the Australian proportion of 94.2%.

Immunisation by provider type for children between the ages of 0 to 6 is shown in Table 6. The proportion of children in the Division who were immunised by a general practitioner was 79.4%, compared to 70.0% for Australia, with 13.9% immunised at a community health centre or by a community health worker, and 5.5% at a local government council.

**Table 6: Childhood immunisation at ages 0 to 6 by provider type, Hunter Urban DGP and Australia, 2003/04**

Provider	Hunter Urban DGP	Australia
	%	%
General practitioner	79.4	70.0
Local government council	5.5	16.6
Community health centre/ worker	13.9	9.8
Public hospital	1.2	2.1
Aboriginal health service/ worker	0.0	0.9
Other*	0.0	0.6
<b>Total: Per cent</b>	<b>100.0</b>	<b>100.0</b>
<b>Number</b>	<b>83,434</b>	<b>3,843,610</b>

\* Includes immunisations in/ by State Health Departments, RFDS and private hospitals

<sup>2</sup>The FWE value is calculated for each GP location by dividing the GP's total Medicare billing (Schedule fee value of services provided during the reference period) by the mean billing of full-time doctors in that derived major speciality for the reference period. Thus, a GP earning 20% more than the mean billing of full-time doctors is shown as 1.2 FWE: this differs from full-time equivalent (FTE) counts, where the FTE value of any GP cannot exceed 1.0.

## Premature mortality

Deaths at ages below 75 years are used as an indicator of health status, as they largely reflect premature deaths, given the current levels of life expectancy in Australia.

The 'all causes' death rate in the Division at ages 0 to 74 years (328.4 deaths per 100,000 population) is higher than for country New South Wales (318.3) and Australia (290.4): the rates have been age standardised to allow for comparisons between areas, regardless of differences in age profiles between the Division and Australia.


The major causes of premature mortality in the Division, as for country New South Wales and Australia as a whole, are cancer and diseases of the circulatory system (Figure 5). With the exception of stroke, motor vehicle accidents and diabetes mellitus, death rates in the Division for all of the listed conditions and causes are similar to or higher than the rates for country New South Wales and Australia.

The data on which the following chart is based are in Table 12.

**Figure 5: Deaths before 75 years of age by major condition group and selected cause, Hunter Urban DGP‡, country NSW and Australia, 2000-02\***

*Indirectly age standardised rate per 100,000 population*

■ Hunter Urban DGP      ■ Country NSW      ■ Australia



\* 'No.' is the total number of deaths for the 2000-02 period; 'Rate' is an annual rate, based on the 3 year average

‡ See note under 'Data converters and mapping' re calculation of Division totals

## Chronic diseases and risk factors

*The term “chronic disease” describes health problems that persist across time and require some degree of health care management (WHO 2002). Chronic diseases tend to have complex causes, are often long lasting and persistent in their effects, and can produce a range of complications (Thacker et al. 1995). They are responsible for a significant proportion of the burden of disease and illness in Australia and other westernised countries. Given the ageing of the population, this trend is likely to continue.*

*At different life stages, risk factors for chronic diseases and their determinants include genetic predisposition; poor diet and lack of exercise; alcohol misuse and tobacco smoking; poor intra-uterine conditions; stress, violence and traumatic experiences; and inadequate living environments that fail to promote healthy lifestyles (NPHP 2001). Risk factors are also more prevalent in areas of low socioeconomic status, and in communities characterised by low levels of educational attainment; high levels of unemployment; substantial levels of discrimination, interpersonal violence and exclusion; and poverty. There is a higher prevalence of risk factors among Indigenous communities, and other socioeconomically disadvantaged Australians (NPHP 2001).*

### Background

In this section, estimates of the prevalence of selected chronic diseases and risk factors, and two summary measures of health, are shown for the Division‡, and for SLAs within the Division: note that the estimates have been predicted from self-reported data, and are not based on clinical records or physical measures. The chronic diseases and risk factors are those for which sufficiently reliable estimates can be made for the Division from national survey data. The process by which the estimates have been made, and details of their limitations, are described in the Notes section, pages 15-16. The data on which the following charts are based are in Table 13.

The estimates provide information of relevance to a number of the National Health Priority Areas (NHPAs – asthma; cardiovascular health; diabetes mellitus; injury prevention and control; mental health; and arthritis and musculoskeletal conditions: estimates have not been made for cancer control, the other NHPA). The risk factors for which estimates have been made are those which are accepted as being associated with these important chronic conditions. They are overweight (not obese), obesity, smoking, lack of exercise and high-risk alcohol use.

*The numbers are estimates for an area, not measured events as are death statistics: they should be used as indicators of likely levels (and not actual levels) of a condition or risk factor in an area.*

### Prevalence estimates: chronic disease‡

It is estimated that, with the exceptions of respiratory system diseases, musculoskeletal system diseases and osteoporosis (females), relatively more people in Hunter Urban DGP reported having any of the selected chronic conditions than in Australia as a whole (Figure 6): that is, the prevalence rates per 1,000 population were higher.

### Prevalence estimates: self-reported health‡

The NHS includes two measures of self-reported health. One is the Kessler Psychological Distress Scale–10 items (K–10). This is a scale of non-specific psychological distress based on 10 questions about negative emotional states in the four weeks prior to interview, asked of respondents 18 years and over (ABS 2002). The other asks respondents aged 15 years and over to rate their health on a scale from ‘excellent’, through ‘very good’, ‘good’ and ‘fair’, to ‘poor’ health.


The population of the Division aged 18 years and over is estimated to have marginally more people with very high psychological distress levels as measured by the K–10 (Figure 7) compared to Australia. The proportion of the population aged 15 years and over estimated to have reported their health as ‘fair’ or ‘poor’ is also above the national average.

‡ See note under ‘Data converters and mapping’ re calculation of Division totals

**Figure 6: Estimates\* of chronic disease and injury, Hunter Urban DGP‡, country New South Wales and Australia, 2001**

*Indirectly age standardised rate per 1,000 population*

■ Hunter Urban DGP      □ Country NSW      ■ Australia




\* 'No.' is a weighted estimate of the number of people in Hunter Urban DGP reporting each chronic condition and is derived from synthetic predictions from the 2001 NHS

‡ See note under 'Data converters and mapping' re calculation of Division totals

**Figure 7: Estimates\* of measures of self-reported health, Hunter Urban DGP‡, country New South Wales and Australia, 2001**

*Indirectly age standardised rate per 1,000 population*

■ Hunter Urban DGP      □ Country NSW      ■ Australia



\* 'No.' is a weighted estimate of the number of people in Hunter Urban DGP reporting under these measures and is derived from synthetic predictions from the 2001 NHS


<sup>1</sup> Kessler 10

‡ See note under 'Data converters and mapping' re calculation of Division totals

## Prevalence estimates: risk factors‡

The Hunter Urban DGP had relatively higher reported rates (when compared to the Australian population) for all of the selected risk factors (Figure 8)

**Figure 8: Estimates\* of selected risk factors, Hunter Urban DGP‡, country New South Wales and Australia, 2001**




\* 'No.' is a weighted estimate of the number of people in Hunter Urban DGP with these risk factors and has been predicted using data from the 2001 NHS and known data for the Division

‡ See note under 'Data converters and mapping' re calculation of Division totals

The following maps provide details of the geographic distribution, at the SLA level, of the estimated prevalence of chronic disease (Map 2), self-reported health (Map 3) and risk factors associated with chronic disease (Map 4).

In the following maps, users should note that the estimates shown for part SLAs in the Division (see Table 11, page 18, for the per cent of SLA population in the Division) represent the estimates for the whole SLA, and not just the part shown. However, SLAs with only a small proportion of their population in the Division are likely to have little influence on the total estimates for the Division, which have been based on the percentage of the SLA population in the Division.

Map 2: Estimates\* of chronic disease and injury by SLA, Hunter Urban DGP, 2001



\* The estimates are synthetic predictions of the prevalence of these conditions: see Notes on the data.




<sup>#</sup> Data not mapped: see Notes on the data.

Map 3: Estimates\* of measures of self-reported health by SLA, Hunter Urban DGP, 2001

Very high psychological distress levels [K-10<sup>1</sup>] (18+ years)

Fair or poor self-assessed health status (15+ years)

Per cent difference from Australian average



<sup>1</sup> Kessler 10

\* The estimates are synthetic predictions of the prevalence of these conditions: see Notes on the data.




# Data not mapped: see Notes on the data.

Map 4: Estimates\* of selected risk factors by SLA, Hunter Urban DGP, 2001

Overweight (not obese) males (15+ years)

Obese males (15+ years)




Overweight (not obese) females (15+ years)



Obese females (15+ years)



Smokers (18+ years)

Physical inactivity (15+ years)



High health risk due to alcohol consumed (18+ years)

Per cent difference from Australian average



\* The estimates are synthetic predictions of the prevalence of these conditions: see Notes on the data.

# Data not mapped: see Notes on the data.

# Notes on the data

## Data sources and limitations

### General

References to 'country New South Wales' relate to New South Wales excluding the Sydney Statistical Division.

### Data sources

Table 7 details the data sources for the material presented in this profile.

**Table 7: Data sources**

Section	Source
<b>Key indicators</b>	
GP services per head of population	GP services data supplied by Department of Health and Ageing, 2003/04 Population data: Estimated Resident Population, ABS, mean of 30 June 2003 and 30 June 2004 populations
<b>Socio-demographic profile</b>	
Figures 1 and 2; Table 1	Estimated Resident Population, ABS, 30 June for the periods shown
Tables 2, 3 and 4; Figures 3 and 4	Data were extracted by postal area from the ABS Population Census 2001 <sup>1</sup> , except for the following indicators: - <i>Indigenous</i> – Experimental estimates of Aboriginal and Torres Strait Islander people, ABS 2001 (unpublished) - <i>Full-time secondary education participation at age 16</i> – Census 2001 (unpublished) - <i>Households receiving rent assistance</i> – Centrelink, December Quarter 2001 (unpublished) - <i>Unemployment rate / Labour force participation</i> – extracted from <i>Small Area Labour Markets Australia</i> , June Quarter 2003, Department of Employment and Workplace Relations
Map 1; Table 9	ABS SEIFA package, Census 2001
<b>General medical practitioner (GP) supply</b>	
Table 5	GP data supplied by Department of Health and Ageing, 2003/04 Population estimates used in calculating the population per GP rates are the: - Census count <sup>2</sup> , ABS Population Census 2001, scaled to 2003/04 - Usual Resident Population <sup>3</sup> , ABS Population Census 2001, scaled to 2003/04 - Day-time population: calculated from journey to work data, ABS Population Census (URP) 2001 (unpublished); and 2001 Census URP, scaled to 2003/04 - Estimated Resident Population, ABS, June 2003/2004
<b>Immunisation</b>	
Text comment: 1 year olds	National Centre for Immunisation Research and Surveillance, 2002
Table 6	Australian Childhood Immunisation Register, Health Insurance Commission, 2003/04 (unpublished)
<b>Premature mortality</b>	
Figure 5; Table 12	ABS Deaths, 2000 to 2002
<b>Chronic diseases and associated risk factors<sup>4</sup></b>	
Figures 6, 7 and 8; Maps 2, 3 and 4; Table 13	Estimated from 2001 National Health Survey (NHS), ABS (unpublished)

<sup>1</sup> All data extracted from Usual Residents Profile, except for data variables only released in the Basic Community Profile

<sup>2</sup> *Census count* - those counted in the Division on Census night, including tourists, business people and other visitors

<sup>3</sup> *Usual Resident Population* - those who usually live there and who were in Australia at the time and would have provided details in the Census at the address where they were counted

<sup>4</sup> See notes below

## Chronic diseases and associated risk factors

The data for chronic conditions and risk factors for SLAs have been estimated from the 2001 National Health Survey (NHS), conducted by the ABS: see note below on synthetic estimates. The NHS sample includes the majority of people living in private households, but excludes the most remote areas of Australia. These areas cover 86.4% of Australia's land mass and comprise just 3% of the total population, however, 28% of Australia's Indigenous population live in these areas. Thus it has not been possible to produce these estimates for Divisions with relatively high proportions of their population in the most remote areas of Australia.

The data for chronic conditions and risk factors are self-reported data, reported to interviewers in the 2001 NHS. Table 8 includes notes relevant to this data.

**Table 8: Notes on estimates of chronic diseases and associated risk factors**

Indicator	Notes on the data
<b>Estimates of chronic disease and injury</b> (Figure 6 and Map 2)	
Long term conditions	- Respondents were asked whether they had been diagnosed with any long term health condition (a condition which has lasted or is expected to last for 6 months or more), and were also asked whether they had been told by a doctor or nurse that they had asthma, cancer, heart and circulatory conditions, and/or diabetes
Injury event	- Injuries which occurred in the four weeks prior to interview
<b>Estimates of measures of self-reported health</b> (Figure 7 and Map 3)	
Very high psychological distress levels (K10)	- Derived from the Kessler Psychological Distress Scale-10 items (K-10), which is a scale of non-specific psychological distress based on 10 questions about negative emotional states in the 4 weeks prior to interview. 'Very high' distress is the highest level of distress category (of a total of four categories)
Fair or poor self-assessed health status	- Respondent's general assessment of their own health, against a five point scale from excellent through to poor – 'fair' or 'poor' being the two lowest in the scale
<b>Estimates of selected risk factors</b> (Figure 8 and Map 4)	
Overweight (not obese)	- Based on self-reported height and weight; BMI calculated and grouped into categories (to allow reporting against both WHO and NHMRC guidelines) - overweight: 25.0 to less than 30.0
Obese	- Based on self-reported height and weight; BMI calculated and grouped into categories (to allow reporting against both WHO and NHMRC guidelines) – obese: 30.0 and greater
Smokers	- Respondent's undertaking regular (or daily) smoking at the time of interview
Physical inactivity	- Did not exercise in the two weeks prior to interview through sport, recreation or fitness (including walking) – excludes incidental exercise undertaken for other reasons, such as for work or while engaged in domestic duties
High health risk due to alcohol consumed	- Respondent's estimated average daily alcohol consumption in the seven days prior to interview (based on number of days and quantity consumed). Alcohol risk levels were grouped according to NHMRC risk levels for harm in the long term, with 'high risk' defined as a daily consumption of more than 75 ml for males and 50 ml for females

**Note:** For a full description, refer to *ABS 2001 National Health Survey, Cat. No. 4364.0* and *ABS 2001 Health Risk Factors, Cat. No. 4812.0*

## Methods

### Synthetic estimates

The estimates of the prevalence of chronic disease and associated risk factors have been predicted for a majority of SLAs across Australia, using modelled survey data collected in the 2001 ABS National Health Survey (NHS) and known characteristics of the area. A synthetic prediction can be interpreted as the likely value for a 'typical' area with those characteristics: the SLA is the area level of interest for this project (where SLAs had small populations they were grouped to larger areas). This work was undertaken by the Australian Bureau of Statistics, as they hold the NHS unit record files: the small area data were compiled by PHIDU.

The approach used is to undertake an analysis of the survey data for Australia to identify associations in the NHS data between the variables that we wish to predict at the area level (eg. prevalence of chronic conditions and risk factors) and the data we have at the area level (eg. socioeconomic status, use of health services). The relationship between these variables for which we have area level data (the predictors) and the reporting of chronic conditions in the NHS is also a part of the model that is developed by the ABS. For example, such associations might be between the number of people reporting specified chronic conditions in the NHS and:

- the number of hospital admissions (in total, to public and to private hospitals, by age, sex and diagnosis),
- socioeconomic status (as indicated by Census data, or for recipients of government pensions and benefits), and
- the number of visits to a general medical practitioner.

The results of the modelling exercise are then applied to the SLA counts of the predictors. The prediction is, effectively, the likely value for a typical area with those characteristics. The raw numbers were then age-standardised, to control for the effects of differences in the age profiles of areas.

*The numbers are estimates for an area, not measured events as are death statistics: they should be used as indicators of likely levels of a condition or risk factor in an area.*

### Premature deaths

Details of deaths by SLA were purchased from the ABS. The raw numbers were then age-standardised, by the indirect method, to control for the effects of differences in the age profiles of areas.

### Data converters and mapping

#### Conversion to Division of data available by postcode

The allocation of postcodes to Divisions was undertaken using information from the Department of Health and Ageing's web site, which shows the proportion of a postcode in a Division (Table 10).

#### Conversion to Division of data available by SLA

(marked in this profile as ‡ See note under 'Data converters and mapping' re calculation of Division total)

Where the data presented in these profiles were only available by SLA they have been converted to Division of General Practice areas using a concordance based on data at the 2001 Census. A copy of the concordance is included in the Population data: A Guide for Divisions of General Practice: it is also available from the Divisions' data area on PHIDU web site.

In brief, the concordance splits the data (eg number of deaths) for each SLA across one or more Divisions. The proportion of an SLA's data that is allocated to each Division was calculated from (a) CD level Census 2001 data that splits SLAs across approximations to postcodes (referred to as postal areas) and (b) data on the DoHA website that splits postcodes across Divisions. This concordance can be adjusted to meet any new configuration of Division boundaries based on the 2001 Collection Districts, or combinations thereof.

The estimated population of each SLA in this Division is shown in Table 11.

#### Mapping

In some Divisions the maps may include a very small part of an SLA which has not been allocated any population, or either has a population of less than 100 or has less than 1% of the SLA's total population: these areas are mapped with a pattern.

## Supporting information

This and other information is also available at [www.publichealth.gov.au](http://www.publichealth.gov.au)

### A definition of population health

Population health, in the context of general practice, has been defined<sup>1</sup> as:

*“The prevention of illness, injury and disability, reduction in the burden of illness and rehabilitation of those with a chronic disease. This recognises the social, cultural and political determinants of health. This is achieved through the organised and systematic responses to improve, protect and restore the health of populations and individuals. This includes both opportunistic and planned interventions in the general practice setting.”*

The key determinants of health are social support networks, employment and working conditions, social environments, physical environments, geographical isolation, personal health practices, healthy child development, ageing and disability, biology and genetic endowment, health services, gender and culture.

In the Aboriginal and Torres Strait Islander context this means that a population health approach to health services will assist in ensuring “that Aboriginal and Torres Strait Islander people enjoy a healthy life equal to that of the general population, that is enshrined by a strong living culture, dignity and justice”.<sup>2</sup> This recognises the importance of achieving improvements to Aboriginal and Torres Strait Islander health and respects the particular health issues facing Indigenous people.

<sup>1</sup> “The role of general practice in population health – A Joint Consensus Statement of the General Practice Partnership Advisory Council and the National Public Health Partnership Group” (Joint Advisory Group on General Practice and Population Health 2001)

<sup>2</sup> As defined in the Strategic Framework for Aboriginal and Torres Strait Islander Health

### SEIFA scores

Following the 2001 Census, the Australian Bureau of Statistics (ABS) produced four socioeconomic indexes for areas (SEIFA). The indexes describe various aspects of the socioeconomic make-up of populations in areas, using data collected in the 2001 Census. The Index of Relative Socio-Economic Disadvantage (labelled ‘Disadvantage’ in Table 9) includes all variables that either reflect or measure disadvantage. The Index of Advantage/Disadvantage is used to rank areas in terms of both advantage and disadvantage: any information on advantaged persons in an area will offset information on disadvantaged persons in the area. The Index of Economic Resources and the Index of Education and Occupation were targeted towards specific aspects of advantage/disadvantage.

For further information on the composition and calculation of these indexes see the ABS Information Paper ABS Cat No. 2039.0 available on the ABS web site [www.abs.gov.au](http://www.abs.gov.au). The scores for these indexes for each Statistical Local Area (SLA) or part SLA in Hunter Urban DGP are shown in Table 9.

**In using this table, users should note that the index score shown for SLAs with less than 100 per cent in the Division represents the score for the whole SLA, and not just the part shown. However, SLAs with small proportions may have little influence on the average index score for the Division which has been based on the postcodes in the Division.**

**Table 9: SEIFA scores by SLA, Hunter Urban DGP, 2001**

SLA code	SLA name (& per cent of SLA in the Division)	Index score			
		Disadvantage	Advantage	Economic Resources	Education & Occupation
11720	Cessnock (13.7)	926	904	932	890
12700	Dungog (18.2)	991	948	936	956
13400	Great Lakes (10.3)	967	918	906	940
14650	Lake Macquarie (98.6)	985	971	971	968
15050	Maitland (98.6)	962	958	974	944
15901	Newcastle - Inner (100.0)	1022	1098	1060	1147
15902	Newcastle - Remainder (100.0)	970	981	968	997
16400	Port Stephens (59.7)	974	952	960	948
17000	Singleton (6.1)	986	985	1030	938
18550	Wyong (1.6)	959	938	958	932

\* Proportions are approximate and are known to be incorrect in some cases, due to errors in the concordance used to allocate CDs to form postal areas

## Statistical geography of the Hunter Urban DGP

The Hunter Urban DGP covers 3,050 square kilometres, based on 2001 SLA data.

The postcodes in the Division (as per the Department of Health and Ageing website) are shown below (Table 10).

**Table 10: Postcodes in Hunter Urban DGP, 2004**

Postcode	Per cent of postcode population in the Division*	Postcode	Per cent of postcode population in the Division*	Postcode	Per cent of postcode population in the Division*
2259	5	2292	100	2308	100
2264	100	2293	100	2310	100
2265	100	2294	100	2314	100
2267	100	2295	100	2318	100
2278	100	2296	100	2319	100
2280	100	2297	100	2320	100
2281	100	2298	100	2321	85
2282	100	2299	100	2322	100
2283	100	2300	100	2323	99
2284	100	2301	67	2324	78
2285	100	2302	100	2325	2
2286	100	2303	100	2326	5
2287	100	2304	100	2330	2
2289	100	2305	100	2334	84
2290	100	2306	100	2335	51
2291	100	2307	100		

\* Proportions are approximate

Source: Department of Health and Ageing web site (accessed online version as at February 2005):

<http://www.health.gov.au/internet/wcms/publishing.nsf/Content/health-pcd-programs-divisions-divspsc.htm>

Statistical Local Areas (SLAs) are defined by the Australian Bureau of Statistics to produce areas for the presentation and analysis of data. In country New South Wales, SLAs are of the same size or, generally, smaller than Local Government Areas (LGAs). In this Division, one Local Government Area (LGA), Newcastle, has been split into two SLAs, Inner and Remainder. All of these SLAs, and all or parts of the other SLAs listed in Table 11 comprise the Division.

**Table 11: SLAs in the Hunter Urban DGP by 2001 boundaries**

SLA code	SLA name	Per cent of the SLA's population in the Division*	Estimate of the SLA's 2004 population in the Division
11720	Cessnock	13.7	6,563
12700	Dungog	18.2	1,520
13400	Great Lakes	10.3	3,519
14650	Lake Macquarie	98.6	186,346
15050	Maitland	98.6	59,017
15901	Newcastle - Inner	100.0	6,172
15902	Newcastle - Remainder	100.0	139,180
16400	Port Stephens	59.7	37,212
17000	Singleton	6.1	1,332
18550	Wyong	1.6	2,227

\* Proportions are approximate and are known to be incorrect in some cases, due to errors in the concordance used to allocate CDs to form postal areas

## Supporting data

The data used in Figure 5 to illustrate the rates of premature mortality in the Division are shown below in Table 12.

**Table 12: Deaths before 75 years of age by major condition group and selected cause, Hunter Urban DGP‡, country New South Wales and Australia, 2000-02\***

*Indirectly age standardised rate per 100,000 population*

Variable	Hunter Urban DGP‡		Country NSW		Australia	
	No.	Rate	No.	Rate	No.	Rate
<b>Circulatory system diseases</b>	1,063	82.8	6,468	83.4	38,357	72.3
Ischaemic heart disease	690	53.7	3,929	50.6	23,364	44.1
Cerebrovascular disease – stroke	160	12.4	1,080	13.8	6,920	13.0
<b>Cancer</b>	1,633	129.1	9,113	119.2	60,603	114.3
Cancer of the trachea, bronchus & lung	338	26.4	1,980	25.4	12,715	24.0
<b>Respiratory system diseases</b>	282	21.7	1,700	21.7	9,726	18.3
Chronic lower respiratory disease	200	15.3	1,209	15.3	6,657	12.6
<b>Injuries and poisonings</b>	435	38.1	2,541	39.5	18,573	35.0
Suicide	175	15.5	888	14.0	6,706	12.6
Motor vehicle accidents	107	9.4	809	12.7	5,014	9.5
<b>Other causes</b>	695	56.5	3,998	54.6	26,735	50.4
Diabetes mellitus	83	6.4	442	9.4	3,734	7.0

\* 'No.' is the total number of deaths for the 2000-02 period; 'Rate' is an annual rate, based on the 3 year average

‡ See note under 'Data converters and mapping' re calculation of Division totals

The rates used to illustrate the prevalence estimates of chronic disease and injury (Figure 6), measures of self-reported health (Figure 7), and selected risk factors (Figure 8), are shown in Table 13 below.

**Table 13: Estimates of chronic diseases and associated risk factors, Hunter Urban DGP‡, country New South Wales and Australia, 2001**

*Indirectly age standardised rate per 1,000 population*

Variable	Hunter Urban DGP‡	Country NSW	Australia
<b>Chronic disease and injury (Figure 6)</b>			
Respiratory system diseases	309.8	310.4	310.8
Asthma	128.9	127.9	118.3
Circulatory system diseases	184.0	181.6	171.5
Diabetes type 2	26.2	23.4	23.4
Injury event	125.4	124.0	121.2
Mental & behavioural disorders	109.2	104.3	97.6
Musculoskeletal system diseases	321.5	322.0	326.2
Arthritis	152.2	148.1	138.8
- Osteoarthritis	84.1	81.1	74.9
- Rheumatoid arthritis	25.3	24.8	23.6
Osteoporosis (females)	24.0	24.1	26.4
<b>Measures of self-reported health (Figure 7)</b>			
Very high psychological distress levels (18+ years)	39.6	38.9	36.6
Fair or poor self-assessed health status (15+ years)	198.2	189.5	184.0
<b>Risk factors (Figure 8)</b>			
Overweight (not obese) males (15+ years)	399.6	397.0	389.7
Obese males (15+ years)	169.5	167.5	145.9
Overweight (not obese) females (15+ years)	243.7	240.9	223.9
Obese females (15+ years)	161.9	157.5	148.0
Smokers (18+ years)	263.9	269.8	248.0
Physical inactivity (15+ years)	340.4	349.9	315.5
High health risk due to alcohol consumed (18+ years)	45.4	47.4	42.1

‡ See note under 'Data converters and mapping' re calculation of Division totals

## References

Australian Bureau of Statistics (ABS) (2002). *2001 National Health Survey: summary of results*. Australia. (ABS Cat. No. 4364.0). Canberra: ABS.

National Public Health Partnership (NPHP) (2001). *Preventing Chronic Disease: A Strategic Framework*. Melbourne, Victoria.

Thacker S, Stroup D & Rothenberg R (1995). Public health surveillance for chronic conditions: a scientific basis for decisions. *Statistics in Medicine* 14: 629-641.

World Health Organization (2002). *The World Health Report 2002: Reducing Risks, Promoting Healthy Life*. Geneva: World Health Organization.

## Acknowledgements

Funding for these profiles was provided by the Population Health Division of the Department of Health and Ageing (DoHA). Assistance, by way of comment on the profiles and assistance in obtaining some datasets, has also been received from the Primary Care Division of the DoHA, the ABS and the ACIR.

## Further developments and updates

Subject to agreement and funding, a number of developments could be undertaken:

- Details of hospitalisations potentially avoidable through ambulatory care interventions are currently being prepared and will be forwarded to Divisions (and posted on the PHIDU web site) when they are available. Other enhancements will be considered as appropriate datasets become available.

The profiles could be updated as the data are updated. For example:

- Population estimates, avoidable hospitalisations, immunisation, and GP activity and workforce data – annually;
- Chronic disease estimates – three-yearly;
- Census data – five-yearly.

Any developments would be informed by consultation, including with Divisions.

## PHIDU contact details

**For general comments, data issues or enquiries re information on the web site, please contact PHIDU:**

Phone: 08-8303 6236 or e-mail: [PHIDU@publichealth.gov.au](mailto:PHIDU@publichealth.gov.au)