

Appendix 1.1: ICD codes

Table A1 details the ICD-9, and ICD-10 (for Australia)/ ICD-10-AM (for New Zealand) codes for the avoidable mortality causes and the mortality amenable to health care groupings. For this analysis, there were no differences in the relevant codes in the ICD-10 and ICD-10-AM versions.

Table A1: Avoidable mortality and amenable mortality conditions and ICD codes

Age limit: 0 to 74 years, unless otherwise specified

| Major condition group/ condition | ICD-9 | ICD-10 [Aust Codes] / ICD-10-AM [NZ Codes] | Limits (age, sex) | Amenable to health care ¹ |
|--|--|---|----------------------|--|
| Infections | | | | |
| Tuberculosis | 010-018,137 | A15-A19, B90 | | ✓ |
| Selected invasive bacterial and protozoal infections | 034-036, 038, 084, 320, 481, 482, 485, 681, 682 | A38-A41, A46, A48.1 B50-B54, G00, G03, J02.0, J13-J15, J18, L03 | | ✓ |
| Hepatitis | 070 | B15-B19 | | |
| HIV/AIDS | 042 | B20-B24 | | |
| Viral pneumonia and influenza | 480, 487 | J10, J12, J17.1, J21 | | |
| Neoplasms | | | | |
| Lip, oral cavity and pharynx | 140-149 | C00-C14 | | |
| Oesophagus | 150 | C15 | | |
| Stomach | 151 | C16 | | |
| Colorectal | 153, 154 | C18-C21 | | ✓ |
| Liver | 155 | C22 | | |
| Lung | 162 | C33, C34 | | |
| Melanoma of skin | 172 | C43 | | ✓ |
| Nonmelanotic skin | 173 | C44 | | ✓ |
| Breast | 174 | C50 | Female | ✓ |
| Cervix | 180 | C53 | | ✓ |
| Uterus | 179, 182 | C54, C55 | | ✓ |
| Bladder | 188 | C67 | | ✓ |
| Thyroid | 193 | C73 | | ✓ |
| Hodgkin's disease | 201 | C81 | | ✓ |
| Lymphoid leukaemia – acute/chronic | 204.0, 204.1 | C91.0, C91.1 | | ✓ |
| Benign | 210-229 | D10-D36 | | ✓ |
| Nutritional, endocrine and metabolic conditions | | | | |
| Thyroid disorders | 240-246 | E00-E07 | | ✓ |
| Diabetes | 250 | E10-E14 | | ✓ (0.5) |
| Drug use disorders | | | | |
| Alcohol related disease | 291, 303, 305.0, 425.5, 535.3, 571.0-571.3 | F10, I42.6, K29.2, K70 | | |
| Illicit drug use disorders | 292, 304, 305.2-305.9 | F11-F16, F18, F19 | | |

... continued

Table A1: Avoidable mortality and amenable mortality conditions and ICD codes ... continued

Age limit: 0 to 74 years, unless otherwise specified

| Major condition group/ condition | ICD-9 | ICD-10 [Aust Codes] / ICD-10-AM [NZ Codes] | Limits (age, sex) | Amenable to health care ¹ |
|--|-------------------------------------|--|----------------------|--|
| Neurological disorders | | | | |
| Epilepsy | 345 | G40, G41 | | ✓ |
| Cardiovascular diseases | | | | |
| Rheumatic and other valvular heart disease | 390-398 | I01-I09 | | ✓ |
| Hypertensive heart disease | 402 | I11 | | ✓ |
| Ischaemic heart disease | 410-414 | I20-I25 | | ✓ (0.5) |
| Cerebrovascular diseases | 430-438 | I60-I69 | | ✓ (0.5) |
| Aortic aneurysm | 441 | I71 | | |
| Genitourinary disorders | | | | |
| Nephritis and nephrosis | 403, 580-589, 591 | I12, I13, N00-N09, N17- N19 | | ✓ |
| Obstructive uropathy & prostatic hyperplasia | 592, 593.7, 594, 598, 599.6, 600 | N13, N20, N21, N35, N40, N99.1 | | ✓ |
| Respiratory diseases | | | | |
| DVT with pulmonary embolism | 415.1, 451.1 | I26, I80.2 | | |
| COPD | 490-492, 496 | J40-J44 | 45-74 years | |
| Asthma | 493 | J45, J46 | 0-44 years | ✓ |
| Digestive disorders | | | | |
| Peptic ulcer disease | 531-534 | K25-K28 | | ✓ |
| Acute abdomen, appendicitis, intestinal obstruction, cholecystitis/ lithiasis, pancreatitis, hernia | 540-543, 550-553, 574- 577 | K35-K38, K40-K46, K80- K83, K85, K86, K91.5 | | ✓ |
| Chronic liver disease (excluding alcohol related disease) | 571.4-571.9 | K73, K74 | | |
| Maternal & infant causes | | | | |
| Birth defects | 237.70, 740-760 | H31.1, P00, P04, Q00- Q99 | | ✓ |
| Complications of perinatal period | 764-779 | P03, P05-P95 | | ✓ |
| Unintentional injuries | | | | |
| Road traffic injuries | E810-E819 | V01-V04, V06, V09-V80, V87, V89, V99 | | |
| Falls | E880-E886, E888 | W00-W19 | | |
| Fires, burns | E890-E899 | X00-X09 | | |
| Accidental poisonings | E850-E869 | X40-X49 | | |
| Drownings | E910 | W65-W74 | | |
| Intentional injuries | | | | |
| Suicide and self inflicted injuries | E950-E959, E980-E989 | X60-X84, Y87.0, Y10-Y34 | | |
| Violence | E960-E969 | X85-Y09, Y87.1 | | |

¹ Subset list of conditions amenable to health care, denoted as ✓; or ✓ (0.5) to represent 50% of total deaths in the category

Table A2 shows the conditions excluded on the basis that they represented less than 0.1% (rounded at one decimal place) of all deaths, based on an analysis of deaths over a recent three (Australia) or four (New Zealand) year period. Note that when one country met the 0.1 per cent requirement, the condition was retained.

Table A2: Avoidable mortality conditions excluded from analysis¹

| Condition | ICD-9 | ICD-10 (Australia); ICD-10-AM (New Zealand) | Percentage of all deaths (%) | |
|--|---|---|------------------------------|-----------------|
| | | | Aust ² | NZ ² |
| Diarrhoeal disease | 001-009 | A00-A09 | 0.03 | 0.05 |
| Childhood vaccine-preventable diseases | 032-033, 036.0, 037, 041.2, 041.5, 045, 070.2-070.3, 052, 055-056 | A35-A37, A39.0, A49.1, A49.2, A80, B01, B05-B06, B16, J11 | Child only | Child only |
| Sexually transmitted diseases except HIV/AIDS | 090-099, 614.0-614.5, 614.7-616.9, 633 | A50-A64, M02.3, N34.1, N70-N73, N75.0, N75.1, N76.4, N76.6, O00 | 0.01 | 0.01 |
| Testis cancer | 186 | C62 | 0.02 | 0.02 |
| Eye cancer | 190 | C69 | 0.03 | 0.02 |
| Nutritional deficiency anemia | 280-281 | D50-D53 | 0.03 | 0.04 |
| Adrenal disorders | 255.0, 255.4 | E24, E27 | 0.01 | 0.01 |
| Newborn screening disorders | 255.2, 270.1, 271.1 | E25, E70.0, E74.2 | 0.00 | 0.00 |
| Ear infections – Otitis media and mastoiditis | 381-383 | H65-H70 | 0.00 | 0.00 |
| Upper respiratory tract infection | 382-383, 460-465 | J00-J06, H66, H70 | 0.02 | 0.04 |
| Osteomyelitis and other osteopathies of bone | 730 | M86, M89-M90 | 0.02 | 0.03 |
| Complication of pregnancy, labor or the puerperium | 630-632, 634-676 | O01-O99 | 0.01 | 0.01 |
| Sports injuries | E884.0, E886.0, E917.0, E927 | any external cause code V00-Y99 with an activity code of 0 | 0.01 | 0.02 |
| War | E990-E999 | Y36 | 0.00 | 0.00 |
| Total of all deaths | .. | .. | 0.26 | 0.39 |

¹ Condition categories were excluded where they represented less than 0.1 per cent of all deaths in both countries

² Percentages were calculated from total deaths over a three or four year period: for Australia: 1997-99; for NZ: 1996-99

This page intentionally left blank